



West Nile virus disease cases and presumptive viremic blood donors reported to ArboNET, United States, 2004

State	Neuroinvasive disease cases*	Non-neuroinvasive disease cases	Total cases†	Deaths	Presumptive viremic blood donors‡
Arizona	215	176	391	16	36
Arkansas	17	11	28	0	0
California	291	488	779	28	66
Colorado	41	250	291	4	24
Connecticut	0	1	1	0	0
Delaware	0	0	0	0	1
District of Columbia	1	1	2	0	0
Florida	33	8	41	2	3
Georgia	14	7	21	1	4
Idaho	1	2	3	0	0
Illinois	29	31	60	4	1
Indiana	8	5	13	1	1
Iowa	13	10	23	2	4
Kansas	18	25	43	2	4
Kentucky	1	6	7	0	1
Louisiana	85	24	109	7	12
Maryland	10	6	16	0	1
Michigan	13	3	16	0	3
Minnesota	13	21	34	2	2
Mississippi	31	20	51	4	2
Missouri	27	9	36	2	2
Montana	2	4	6	0	0
Nebraska	7	46	53	0	3
Nevada	25	19	44	0	5
New Jersey	1	0	1	0	1
New Mexico	32	56	88	4	15
New York	7	3	10	0	1
North Carolina	3	0	3	0	0
North Dakota	2	18	20	2	1
Ohio	11	1	12	2	0
Oklahoma	16	6	22	3	8
Oregon	0	3	3	0	1
Pennsylvania	10	5	15	2	1
South Carolina	0	2	2	0	0
South Dakota	6	<u>-</u> 45	 51	1	3
Tennessee	13	1	14	0	0
Texas	119	57	176	8	16
Utah	6	5	11	0	0
Virginia	5	0	5	1	0
Wisconsin	5	7	12	2	2
Wyoming	2	8	10	0	0
Total	1,148	1,391	2,539	100	224

^{*}Includes cases reported as meningitis, encephalitis, or acute flaccid paralysis.

[†]Includes confirmed and probable cases.

[‡]Presumptive viremic blood donors (PVDs) are people who had no symptoms at the time of donating blood through a blood collection agency, but whose blood tested positive when screened for the presence of West Nile virus RNA. Some PVDs develop symptoms after donation; of the 224 PVDs, 70 (31%) developed clinical illness and are also included as disease cases.